



LAWRENCE
LIVERMORE
NATIONAL
LABORATORY

Pu and Np Sorption onto Manganese(IV) Oxide Minerals

P. Zhao, M. R. Johnson, S. K. Roberts, M. Zavarin

March 11, 2005

229th ACS National Meeting · San Diego, California, USA
San Diego, CA, United States
March 13, 2005 through March 17, 2005

Disclaimer

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.



Pu and Np Sorption onto Manganese(IV) Oxide Minerals

Pihong Zhao*, Mackenzie R. Johnson**,
Sarah K. Roberts**, Mavrik Zavarin**

*Chemical Biology and Nuclear Science Division


**Environmental Science Division

Lawrence Livermore National Laboratory, 7000 East Avenue, Livermore, CA 94550



Objectives of this Work

- Manganese oxide minerals are present at Yucca Mountain and are likely to present at NTS as well
- These minerals have high sorption capacity
- Manganese sorption data were not included in the radionuclide transport models
- This results in an over-prediction of radionuclide transport




Mineral	Tuff Cone unit Mineral %	Lava Flow Aquifer unit, Mineral %
Chalcedony	24	24
Quartz	5	8
Calcite	3	3
Zeolite	30	35
Clay	11	6
Fe/MnO	20	17
Other	7	7




Experimental Outlines

- Batch sorption Experiments
 - Synthetic minerals birnessite, cryptomelane and pyrolusite
 - K_d vs. pH 3-10,
 - Sorption rate
- Batch desorption experiments
 - sorption reversibility,
 - desorption rate
- Supernatant characterization
 - Solvent extraction
 - Ion Chromatography
 - total carbonate analyses







Manganese(IV) Oxide Minerals



layer structure



tunnel structure



framework structure

Mineral Surface Areas (m^2/g) (BET data)

Birnessite

41.36 +/- 0.08

54.70 +/- 0.13

Cryptomelane

11.79 +/- 0.15






10.54 +/- 0.78

Pyrolusite

2.30 +/- 0.04



SEM Images of three Manganese Oxides Minerals






Birnessite

Cryptomelane

Pyrolusite




Np(V) sorption onto manganese oxides as a function of time



- Similar sorption behavior at equilibrium.
- The solution pH is a primary factor affecting Np sorption.
- Np-carbonate complexation may reduce the sorption at high pH.
- The sorption rate was fast initially and followed by a slow rate over one month.




Np(V) Sorption onto manganese oxides



- The K_d values range from <100 at lower pH to >1000 at higher pH.
- Sorption of Np(V) is comparable when examined in terms of %sorbed or K_d values.
- Sorption per unit surface area: Pyrolusite $>$ Cryptomelane $>$ Birnessite.






Comparison of Np(V) Sorption / Desorption on Birnessite



- Desorption at range pH 3-6 could not be measured
- Desorption was fast and reversible
- Similar desorption data were observed for cryptomelane and pyrolusite.





Pu(IV) sorption onto manganese oxides as a function of time



- Similar sorption behavior at equilibrium regardless minerals.
- pH is a primary factor affecting Pu(IV) sorption.
- Pu-carbonate complexation reduces the sorption at high pH.
- Sorption onto all three minerals were very strong in neutral to weak basic solutions.
- The sorption rate was fast for pH > 7, and was relative slow in weak acidic solution.




Pu(IV) Sorption onto manganese oxides



- The K_d values change in a wide range from $<10 \sim >1E7$.
- K_d values unveil the effect of Pu-carbonate complexation at higher pH.
- Sorption per unit surface area: Pyrolusite > Cryptomelane > Birnessite.



Comparison of Pu(IV) Sorption / Desorption on Birnessite



- Desorption was fast and reversible
- Similar desorption data were observed for cryptomelane and pyrolusite



Np Oxidation States in Supernatants from Desorption Experiment

		Percentage of Np oxidation state		
		4+	5+	6+
Birnessite	pH4	1%	92%	7%
	pH7		80%	20%
	pH10		68%	32%
Cryptomelane	pH4	3%	96%	1%
	pH7	1%	97%	2%
	pH10		99%	1%
Pyrolusite	pH4		98%	2%
	pH7**	---	---	---
	pH10		99%	1%

** Data were not available due to very low Np concentration UCRL-CONF-210424



Pu Oxidation States in Supernatants from Desorption Experiment

		Pu Oxidation States				colloids
		3+	4+	5+	6+	
Birnessite	pH4		1%	66%	28%	5%
	pH7*	10%	40%	10%	40%	
	pH10	1%	55%	7%	37%	
Cryptomelane	pH4			80%	19%	
	pH7**	---	---	---	---	---
	pH10**	---	---	---	---	---
Pyrolusite	pH4		1%	94%	3%	2%
	pH7**	---	---	---	---	---
	pH10		89%	2%	7%	1%

* Data were associated with large errors


** Data were not available due to very low Pu concentration

UCRL-CONF-210424



Summary

- Manganese(IV) oxide minerals are good sorbers for Np(V) and very strong sorbers for Pu(IV).
- Sorption of Np(V) and Pu(IV) was fast, reversible, and pH dependent
- On a mass basis, sorption to all three Mn oxides was similar. Base on the BET data, sorption/m² followed the order: pyrolusite>cryptomelane>birnessite
- The oxidation states of both Pu and Np appeared to change in the desorption,
- Transport of Pu and Np at NTS locations may be significantly affected by the presence of manganese oxide minerals.
- Future modeling efforts should address the role of Mn oxides in radionuclides transport to more accurately predict their transport





Acknowledgements

- Dr. Qinghong Hu
 - Dr. Annie B. Kersting
 - Dr. Timothy P. Rose
-
- Funding for this investigation was provided by the Environment Restoration Division's Underground Test Area Project at the U.S. Department of Energy, National Nuclear Security Administration, Nevada Site Office.
 - This Work was performed under the auspices of the U.S. Department of Energy by UC, Lawrence Livermore National Laboratory under contract W-7405-Eng-48.